


Example Responses – Paper 5

Cambridge IGCSE[™] / IGCSE (9–1) Physics 0625 / 0972

For examination from 2023







Contents

Introduction	4
Question 1	5
Question 2	
Question 3	
Ouestion 4	14

Introduction

The main aim of this booklet is to exemplify standards for those teaching Cambridge IGCSE / IGCSE (9-1) Physics 0625 / 0972.

This booklet contains responses to all questions from June 2023 Paper 51, which have been written by a Cambridge examiner. Responses are accompanied by a brief commentary highlighting common errors and misconceptions where they are relevant.

The question papers and mark schemes are available to download from the School Support Hub

0625 / 0972 June 2023 Question Paper 51 0625 / 0972 June 2023 Mark Scheme 51 0625 / 0972 June 2023 Confidential Instructions Paper 51

Past exam resources and other teaching and learning resources are available from the <u>School Support Hub</u>

1 In this experiment, you will investigate the balancing of a metre ruler.

Carry out the following instructions, referring to Fig. 1.1.




Fig. 1.1

- (a) Place the metre ruler on the pivot at the 50.0 cm mark with its scale facing upwards.
 - Place the object Q with its centre on the metre ruler at the 90.0 cm mark.

Record the distance *y* from the centre of Q to the 100.0 cm end of the ruler.

Examiner comment

A minority of candidates measured the distance on Fig. 1.1 instead of calculating the actual distance.

- **(b)** Place a load P of weight $P = 2.0 \,\text{N}$ on the metre ruler.
 - Adjust the position of P so that the metre ruler is as near as possible to being balanced.
 - (i) Measure, and record in Table 1.1, the distance *x* from the centre of P to the zero end of the ruler. Record the weight *P*. [1]
 - (ii) Repeat the steps above, using loads of weight $P = 3.0 \,\mathrm{N}, 4.0 \,\mathrm{N}, 5.0 \,\mathrm{N}$ and $6.0 \,\mathrm{N}.$

Record all the values of *P* and *x* in Table 1.1. Ensure that the position of object Q on the metre ruler does **not** change.

Table 1.1

P/N	x/cm
2.0	10.2
3.0	23.1
4.0	30.0
5.0	33.8
6.0	36.8


[3]

Examiner comment

While most candidates produced a good set of readings with x values decreasing, some were not awarded the marks for giving the P values to two significant figures or for giving all the x values to the nearest mm.

(c) Plot a graph of P/N (y-axis) against x/cm (x-axis).

Draw the best-fit line.



Examiner comment

The majority of candidates obtained readings that produced a clear curve when plotted. However, some candidates appeared to assume that a straight line graph was expected and so did not gain the fourth graph mark.

(d) Use the graph to find the value of x required to balance the ruler when $P = 3.5 \,\mathrm{N}$.

Show clearly on the graph how you determined the value of *x*.

Examiner comment

Most candidates were successful in this question, but some drew a triangle to work out the gradient of the line. This was not what the question demanded.

2 In this experiment, you will investigate the cooling of water.

Carry out the following instructions, referring to Fig. 2.1.




Fig. 2.1

(a) Use the thermometer to measure the room temperature $\theta_{\rm R}$.

$$\theta_{\mathsf{R}} = 22 \, {}^{\circ}\mathcal{C}$$
 [1]

Examiner comment

Very few candidates gave an unrealistic room temperature.

(b)	(i)	Pour 200 cm ³ of hot water into the beaker. Place the thermometer in the hot water in the
		beaker.

Record in Table 2.1 the temperature θ of the hot water at time t = 0. Immediately start the stop-watch.

Continue recording the temperature in Table 2.1 at 30s intervals until you have seven sets of readings. [2]

(ii) Complete the column headings in Table 2.1.

Table 2.1

	1
t/ s	θ I $^{\circ}C$
0	92
30	84
60	78
90	74
120	71
150	69
180	67

[1]

Examiner comment

- Some candidates gave room temperature as the starting temperature or a temperature between room temperature and the highest recorded temperature due to taking the initial reading too soon.
- A few candidates gave a wrong unit for temperature (e.g. C°). A more common error was to miss the instruction to complete the column headings.
 - (c) (i) Calculate the decrease in temperature $\Delta\theta$ between t=0 and t=180 s.

$$\Delta\theta$$
 =[1]

(ii) Calculate the average rate of cooling *R* of the water using the equation $R = \frac{\Delta \theta}{\Delta t}$, where $\Delta t = 180 \, \text{s}$. Include the unit.

$$R = \frac{0.139 \text{ °C/s}}{1.000 \text{ Colorest}}$$

- (d) A student states that the average rate of cooling of the water decreases as the temperature comes nearer to room temperature.
 - (i) Suggest **one** change to the experiment that you could make to test the statement.

continue to take temperatures for a longer time
[1]

	plot a graph of temperature against time
	[2]
	ner comment pproaches were possible here, all making the same basic point. For example, continuing to take
emperatu	res until close to room temperature or using a higher or lower starting temperature.
(e)	Explain briefly why it is good practice to read the thermometer scale at right angles.
	to avoid parallax error
	[1]

(ii) Suggest how to display the results to make it easier to see the trend in the rate of cooling.

Examiner comment

- Candidates did not need to use the word 'parallax' in their responses. A description of the effect, for example, the value looks different from the other angle, or a reference to avoidance of line-of-sight error, that showed that a candidate was familiar with good practice was enough to be awarded the mark.
- References to avoidance of inaccuracy or human error were too vague to attract the mark.

3 In this experiment, you will investigate the refraction of light using a semicircular transparent block.

Carry out the following instructions using the separate ray-trace sheet provided. You may refer to Fig. 3.1 and Fig. 3.2 for guidance.




Fig. 3.1

- (a) Draw a line across the ray-trace sheet supplied, approximately in the middle. Label the line PQ.
 - Place the transparent block, largest face down, with the straight side on the line **PQ** and the curved side below the line.
 - Draw round the outline of the block. Label the ends of the straight side of the block A
 and B.
 - Remove the block and draw the normal **NL** through the centre of **AB**. Continue the normal so that it passes through the curved side of the block.
 - Label the point C where the normal NL crosses AB.

7

- **(b)** Draw the line **DC** at an angle $i = 30^{\circ}$ to the normal, as shown in Fig. 3.1.
 - Place the paper on the pin board.
 - Place two pins, P₁ and P₂, on line **DC** at a suitable distance apart for this experiment.
 - Replace the block and look from the position of the eye shown in Fig. 3.1 to observe the images of P₁ and P₂ through side **AB** of the block. Adjust your line of sight until the images of P₁ and P₂ appear one behind the other.
 - images of P₁ and P₂ appear one behind the other.

 Place two pins, P₃ and P₄, between your eye and the block so that P₃, P₄, and the images of P₁ and P₂ seen through the block, appear one behind the other.
 - Label the positions of P₁, P₂, P₃ and P₄.
 - Remove the block and the pins.
 - Draw a line joining the positions of P₃ and P₄. Continue the line to AB.
 - Label E, the end of the line furthest from AB.

[3]

(c) Measure the acute angle θ between the line **NL** and the line **CE**. (An acute angle is less than 90°.)

θ =° [2]

Examiner comment

While most candidates constructed the diagram well, many placed the two pins P_1 and P_2 too close together. A minimum distance apart of 5.0 cm was required.

(d) State one precaution that you take to produce an accurate ray trace.

view	the	bases	of the	pins	 	 	 	
				•	 	 	 	
					 	 	 	 [1]

Examiner comment

- Other acceptable responses included 'place pins as far apart as possible', 'use a sharp pencil / draw thin lines' or 'ensure pins are vertical'.
- Some candidates appeared to have learned answers from previous examination paper mark schemes and made suggestions appropriate for different experiments. For example 'use a darkened room' which is appropriate for a lens experiment using an illuminated object.

(e) Place the transparent block on the ray-trace sheet in the position shown in Fig. 3.2.




Fig. 3.2

- Replace pins P_1 and P_2 on line **DC** in the same positions used in **(b)**.
- Observe the images of P_1 and P_2 through the curved side of the block. Adjust your line of sight until the images of P_1 and P_2 appear one behind the other. Place two pins, P_5 and P_6 , between your eye and the block so that P_5 , P_6 , and the images of P_1 and P_2 seen through the block, appear one behind the other.
- Label the positions of P_5 and P_6 .
- Remove the block and the pins.
- Draw a line joining the positions of P_5 and P_6 . Continue the line to **AB**.
- Label F, the end of the line furthest from AB.

[2]

Measure the acute angle θ between the line **NL** and the line **CF**. (An acute angle is less than (f) 90°.)

Examiner comment

Some candidates had difficulty lining up the pins correctly, perhaps due to lack of experience.

4 A student investigates the change in resistance of a lamp filament when the current in the lamp is increased.

The following apparatus is available:

- a power supply
- a low-voltage filament lamp
- · an ammeter
- a voltmeter
- connecting wires.

Other apparatus normally found in a school laboratory is also available.


Plan an experiment to investigate the change in resistance of the lamp filament when the current in the lamp is increased.

Resistance R is given by the equation $R = \frac{V}{I}$, where V is the potential difference (p.d.) across the lamp and I is the current in the lamp.

You are **not** required to do this investigation.

You should:

- draw a diagram of the circuit used
- explain briefly how to do the investigation, including how to change the current
- draw a table, or tables, with column headings, to show how to display your readings (you are not required to enter any readings in the table)
- explain how to use your readings to reach a conclusion.



Method:					
 using the circuit show 	In, measu	ire the cu	rrent in	the lamp and the	
potential difference a	cross the	lamp			
• calculate the resistan	ce				
 repeat with five differ 	rent value	es of curr	ent		
 use a variable resistor 	to chang	ge the cur	rent		
Table:		•			
 write readings and re 	esults in a	table as	shown:		
	1/A	V / V	R/Ω		
					•
Conclusion:					
 plot a graph of resist. 	ance agai	nsi curre	ni.		
				[7	"]

Examiner comment

- Diagram: the main error with the circuit diagram was to place the voltmeter in series with the other components.
- Method: many candidates included unnecessary information in the method, often copying the list of apparatus from the question. Candidates often missed out one of the essential readings (current or potential difference), what to do with the readings (i.e. calculate the resistance) and a method for changing the current.
- Table: the most common error seen in the table was current / I, instead of I / A.
- Conclusion: a clear statement about comparing values of current and resistance as an alternative to the graph
 was acceptable. Instead of a conclusion, some candidates wrote a prediction which was not required and was not
 awarded a mark.