Cambridge International AS & A Level

CANDIDATE
NAME

CENTRE
NUMBER

CANDIDATE
NUMBER

MATHEMATICS **9709/22**

Paper 2 Pure Mathematics 2 **February/March 2022**

1 hour 15 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [ ].

[Turn over

1 Solve the equation $|5x - 2| = |4x + 9|$. [3]

2   A curve has equation $y = 7 + 4\ln(2x + 5)$.

Find the equation of the tangent to the curve at the point $(-2, 7)$, giving your answer in the form $y = mx + c$.   [5]

3 The variables x and y satisfy the equation $y = 3^{2a}a^x$, where a is a constant. The graph of $\ln y$ against x is a straight line with gradient 0.239.

(a) Find the value of a correct to 3 significant figures. [3]

....

....

....

....

....

....

....

....

....

....

....

(b) Hence find the value of x when $y = 36$. Give your answer correct to 3 significant figures. [2]

....

....

....

....

....

....

....

....

....

4 (a) Show that $\sin 2\theta \cot \theta - \cos 2\theta \equiv 1$. [3]

....

....

....

....

....

....

....

....

....

(b) Hence find the exact value of $\sin \frac{1}{6}\pi \cot \frac{1}{12}\pi$. [2]

....

....

....

....

....

....

(c) Find the smallest positive value of θ (in radians) satisfying the equation

$$\sin 2\theta \cot \theta - 3\cos 2\theta = 1.$$ [2]

....

....

....

....

5 **(a)** Given that $y = \tan^2 x$, show that $\dfrac{dy}{dx} = 2\tan x + 2\tan^3 x$. [2]

....

....

....

....

....

....

....

....

....

....

....

....

(b) Find the exact value of $\displaystyle\int_{\frac{1}{4}\pi}^{\frac{1}{3}\pi} (\tan x + \tan^2 x + \tan^3 x)\,dx.$ [6]

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

6 The polynomial p(x) is defined by

$$p(x) = 4x^3 + 16x^2 + 9x - 15.$$

(a) Find the quotient when p(x) is divided by ($2x + 3$), and show that the remainder is -6. [3]

....

....

....

....

....

....

....

....

....

....

....

(b) Find $\displaystyle\int \frac{p(x)}{2x + 3}\, dx$. [2]

....

....

....

....

....

....

....

....

....

(c) Factorise $p(x) + 6$ completely and hence solve the equation

$$p(\operatorname{cosec} 2\theta) + 6 = 0$$

for $0° < \theta < 135°$. [5]

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

7 A curve has equation $e^{2x}y - e^y = 100$.

(a) Show that $\dfrac{dy}{dx} = \dfrac{2e^{2x}y}{e^y - e^{2x}}$. [3]

....

....

....

....

....

....

....

....

....

....

(b) Show that the curve has no stationary points. [2]

....

....

....

....

....

....

....

....

....

....

It is required to find the x-coordinate of P, the point on the curve at which the tangent is parallel to the y-axis.

(c) Show that the x-coordinate of P satisfies the equation

$$x = \ln 10 - \tfrac{1}{2}\ln(2x - 1).$$

[4]

....

....

....

....

....

....

....

....

....

....

(d) Use an iterative formula, based on the equation in part **(c)**, to find the x-coordinate of P correct to 3 significant figures. Use an initial value of 2 and give the result of each iteration to 5 significant figures. [3]

....

....

....

....

....

....

....

....

....

....

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....