



# Cambridge International AS & A Level

CANDIDATE  
NAME


--

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--



**CHEMISTRY**

**9701/52**

Paper 5 Planning, Analysis and Evaluation

**February/March 2024**

**1 hour 15 minutes**

You must answer on the question paper.

No additional materials are needed.

## INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

## INFORMATION


- The total mark for this paper is 30.
- The number of marks for each question or part question is shown in brackets [ ].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **12** pages. Any blank pages are indicated.


- 1 Sea water contains about  $20\text{ g dm}^{-3}$  of chloride ions,  $\text{Cl}^{-}(\text{aq})$ .

The exact concentration of  $\text{Cl}^{-}(\text{aq})$  in sea water can be determined by titration with aqueous silver ions,  $\text{Ag}^{+}(\text{aq})$ , using aqueous potassium chromate(VI),  $\text{K}_2\text{CrO}_4(\text{aq})$ , as an indicator.

When aqueous silver nitrate,  $\text{AgNO}_3(\text{aq})$ , is added to a sample of sea water, silver ions react with chloride ions to form a precipitate of silver chloride.



When all of the  $\text{Cl}^{-}(\text{aq})$  has reacted with  $\text{Ag}^{+}(\text{aq})$ , the presence of unreacted  $\text{Ag}^{+}(\text{aq})$  is detected by chromate(VI) ions,  $\text{CrO}_4^{2-}(\text{aq})$ . A red precipitate of  $\text{Ag}_2\text{CrO}_4(\text{s})$  is seen.



The amount of  $\text{Ag}^{+}(\text{aq})$  reacting with  $\text{Cl}^{-}(\text{aq})$  in the sample of sea water can be calculated in order to determine the concentration of  $\text{Cl}^{-}(\text{aq})$  in the sample of sea water.

A student uses the following method.

- step 1** Use a weighing boat to weigh by difference approximately 10.6 g of  $\text{AgNO}_3(\text{s})$  into a  $100\text{ cm}^3$  glass beaker.
- step 2** Use the sample of  $\text{AgNO}_3(\text{s})$  in the glass beaker to prepare  $250.0\text{ cm}^3$  of  $\text{AgNO}_3(\text{aq})$ .
- step 3** Transfer this solution into a dark brown glass bottle. Label this solution **X**.
- step 4** Collect a sample of sea water and remove any solid material present.
- step 5** Transfer  $10.00\text{ cm}^3$  of the sea water into a conical flask.
- step 6** Add  $1\text{ cm}^3$  of  $\text{K}_2\text{CrO}_4(\text{aq})$  to the conical flask.
- step 7** Rinse a burette in preparation for the titration.
- step 8** Fill the burette with solution **X**.
- step 9** Slowly add solution **X** to the conical flask until the white precipitate turns red. This is the end-point.

- (a) Describe how the student should carry out step 1. Include a table in your answer to show how this process is recorded.

.....  
 .....

[2]

- (b) Describe how the student should prepare 250.0 cm<sup>3</sup> of AgNO<sub>3</sub>(aq) in step 2, starting with the AgNO<sub>3</sub>(s) in the 100 cm<sup>3</sup> beaker in step 1.

.....  
 .....

[3]

- (c) Suggest why solution X is kept in a dark brown glass bottle in step 3 rather than a colourless glass bottle.

..... [1]

- (d) Suggest how solid material should be removed from sea water in step 4.

..... [1]

- (e) Identify the most appropriate piece of equipment that you would use to:

- (i) transfer 10.00 cm<sup>3</sup> of sea water from the dark brown bottle to a conical flask in step 5

..... [1]

- (ii) add 1 cm<sup>3</sup> of K<sub>2</sub>CrO<sub>4</sub>(aq) to the conical flask in step 6.

..... [1]

- (f) Chromate(VI) solutions are known to be carcinogenic. State what precaution should be taken when using K<sub>2</sub>CrO<sub>4</sub>(aq) in step 6 other than wearing safety goggles.

..... [1]

- (g) State what the burette should be rinsed with in step 7.

..... [1]

(h) The student obtains the results shown in Table 1.1.

**Table 1.1**

	rough titration	titration 1	titration 2	titration 3
final volume / cm <sup>3</sup>	23.40	45.75	22.60	45.05
initial volume / cm <sup>3</sup>	0.00	23.40	0.00	22.60
titre / cm <sup>3</sup>				

(i) Complete Table 1.1. [1]

(ii) Calculate the mean titre to be used in the calculations. Show your working.

mean titre = ..... cm<sup>3</sup> [1]

(iii) Use the mean titre from (h)(ii) to calculate the concentration of chloride ions in the sample of sea water.

Assume the mass of solid silver nitrate used in step 2 was 10.62 g.

concentration = ..... mol dm<sup>-3</sup> [3]

(iv) Calculate the percentage error in the titre in titration 2.  
Show your working.

percentage error = ..... % [1]


(i) Spectroscopic analysis of the sample of sea water accurately determined the concentration of Cl<sup>-</sup>(aq) to be lower than that determined by titration with Ag<sup>+</sup>(aq).

Suggest why the student's method gave a higher value.

.....  
..... [1]

[Total: 18]

- 2 A student wants to investigate the rate of the hydrolysis of methyl methanoate,  $\text{HCOOCH}_3$ .



The reaction is catalysed by dilute hydrochloric acid,  $\text{HCl}(\text{aq})$ .

The amount of methanoic acid,  $\text{HCOOH}$ , produced as the reaction progresses can be monitored by titration with aqueous sodium hydroxide,  $\text{NaOH}(\text{aq})$ , of known concentration using thymolphthalein as the indicator.

To determine this, the volume of  $\text{NaOH}(\text{aq})$  needed to neutralise the  $\text{H}^+(\text{aq})$  from the catalyst needs to be found beforehand.

The student uses the following procedure.

**step 1** Add approximately  $150\text{ cm}^3$  of iced water to a  $250\text{ cm}^3$  conical flask, **A**.

**step 2** Add  $200\text{ cm}^3$  of  $0.250\text{ mol dm}^{-3}$   $\text{HCl}(\text{aq})$  to a  $500\text{ cm}^3$  conical flask, **B**.

Conical flask **B** is the flask in which the reaction takes place.

**step 3** Transfer  $2.00\text{ cm}^3$  of  $0.250\text{ mol dm}^{-3}$   $\text{HCl}(\text{aq})$  from conical flask **B** to conical flask **A**. Carry out a single titration of the contents of conical flask **A** with  $\text{NaOH}(\text{aq})$  of known concentration.

**step 4** Add  $10.0\text{ cm}^3$  of methyl methanoate to conical flask **B**, swirl the reaction mixture and immediately start a stopwatch.

**step 5** After 1 minute transfer  $2.00\text{ cm}^3$  of the reaction mixture from conical flask **B** into conical flask **A**. Carry out a further single titration of the contents of conical flask **A** against  $\text{NaOH}(\text{aq})$ . Do **not** empty the contents of conical flask **A** between titrations.

**step 6** After 10 minutes transfer  $2.00\text{ cm}^3$  of the reaction mixture from conical flask **B** into conical flask **A**. Titrate the contents of conical flask **A** against  $\text{NaOH}(\text{aq})$ .

**step 7** Repeat step 6 at intervals of 10 minutes for 1 hour.

- (a) State which step is used to determine the concentration of  $\text{H}^+(\text{aq})$  ions from the catalyst in the mixture.

..... [1]

- (b) The iced water in conical flask **A** is used to significantly reduce the rate of reaction.

Suggest **two** reasons why the rate of reaction is significantly reduced when the reaction mixture is transferred to conical flask **A**.

reason 1 .....

.....

reason 2 .....

.....

[2]

(c) Table 2.1 shows the readings taken by the student.

The titrations in steps 4–7 show the volume of NaOH(aq) needed to neutralise both the  $\text{H}^+(\text{aq})$  ions from the catalyst,  $\text{HCl}(\text{aq})$ , and from the  $\text{HCOOH}$  produced in the reaction.

volume of NaOH(aq) needed, in  $\text{cm}^3$ , to neutralise  $\text{H}^+(\text{aq})$  from catalyst =  $11.40 \text{ cm}^3$

volume of NaOH(aq), in  $\text{cm}^3$ , used to neutralise  $\text{H}^+(\text{aq})$  from  $\text{HCOOH}$  at time,  $t = V_t$

volume of NaOH(aq), in  $\text{cm}^3$ , used to neutralise  $\text{H}^+(\text{aq})$  from  $\text{HCOOH}$  at 60 min =  $V_\infty$

**Table 2.1**

reading	time, $t$ /min	total volume of NaOH(aq) needed to neutralise total amount of $\text{H}^+(\text{aq})$ / $\text{cm}^3$	$V_t$ / $\text{cm}^3$	$(V_\infty - V_t)$ / $\text{cm}^3$
1	1	12.60		
2	13	17.70		
3	20	19.90		
4	30	22.10		
5	40			
6	50	24.90		
7	60	25.90		

The student forgot to take reading 5.

(i) Complete Table 2.1. [2]

(ii) Identify the independent variable.

..... [1]

(iii) Identify **one** variable that needs to be controlled, apart from concentrations and volumes of solutions.


..... [1]

(iv) Reading 2 should have been taken at 10 minutes and **not** at 13 minutes.

State whether this result should have been included or not. Explain your answer.

.....  
 ..... [1]

- (v) Plot a graph on the grid in Fig. 2.1 to show the relationship between  $(V_{\infty} - V_t)$  and time. Use a cross ( $\times$ ) to plot each data point. Draw a line of best fit.



**Fig. 2.1**

[2]

- (vi) Reading 5 was **not** taken. Use the graph to predict the total volume of NaOH(aq) needed to neutralise the total amount of  $H^+$ (aq) at 40 minutes.

volume of NaOH(aq) = ..... [1]

- (vii) It is **not** possible to repeat the experiment.

State whether the data from the experiment is reliable. Justify your answer.

.....  
 ..... [1]

[Total: 12]









**Important values, constants and standards**

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 $\text{J g}^{-1} \text{ K}^{-1}$ )

## The Periodic Table of Elements

Group																			
1	2																	17	18
		<div style="display: flex; justify-content: space-between;"> <div style="border: 1px solid black; padding: 2px;">1 H hydrogen 1.0</div> <div style="border: 1px solid black; padding: 2px;">2 He helium 4.0</div> </div>																	
		<div style="display: flex; justify-content: space-between;"> <div style="border: 1px solid black; padding: 2px;">3 Li lithium 6.9</div> <div style="border: 1px solid black; padding: 2px;">4 Be beryllium 9.0</div> </div>																	
		<div style="display: flex; justify-content: space-between;"> <div style="border: 1px solid black; padding: 2px;">5 B boron 10.8</div> <div style="border: 1px solid black; padding: 2px;">6 C carbon 12.0</div> <div style="border: 1px solid black; padding: 2px;">7 N nitrogen 14.0</div> <div style="border: 1px solid black; padding: 2px;">8 O oxygen 16.0</div> <div style="border: 1px solid black; padding: 2px;">9 F fluorine 19.0</div> <div style="border: 1px solid black; padding: 2px;">10 Ne neon 20.2</div> </div>																	
		<div style="display: flex; justify-content: space-between;"> <div style="border: 1px solid black; padding: 2px;">11 Na sodium 23.0</div> <div style="border: 1px solid black; padding: 2px;">12 Mg magnesium 24.3</div> </div>																	
		<div style="display: flex; justify-content: space-between;"> <div style="border: 1px solid black; padding: 2px;">13 Al aluminium 27.0</div> <div style="border: 1px solid black; padding: 2px;">14 Si silicon 28.1</div> <div style="border: 1px solid black; padding: 2px;">15 P phosphorus 31.0</div> <div style="border: 1px solid black; padding: 2px;">16 S sulfur 32.1</div> <div style="border: 1px solid black; padding: 2px;">17 Cl chlorine 35.5</div> <div style="border: 1px solid black; padding: 2px;">18 Ar argon 39.9</div> </div>																	
		<div style="display: flex; justify-content: space-between;"> <div style="border: 1px solid black; padding: 2px;">19 K potassium 39.1</div> <div style="border: 1px solid black; padding: 2px;">20 Ca calcium 40.1</div> </div>																	
		<div style="display: flex; justify-content: space-between;"> <div style="border: 1px solid black; padding: 2px;">21 Sc scandium 45.0</div> <div style="border: 1px solid black; padding: 2px;">22 Ti titanium 47.9</div> <div style="border: 1px solid black; padding: 2px;">23 V vanadium 50.9</div> <div style="border: 1px solid black; padding: 2px;">24 Cr chromium 52.0</div> <div style="border: 1px solid black; padding: 2px;">25 Mn manganese 54.9</div> <div style="border: 1px solid black; padding: 2px;">26 Fe iron 55.8</div> <div style="border: 1px solid black; padding: 2px;">27 Co cobalt 58.9</div> <div style="border: 1px solid black; padding: 2px;">28 Ni nickel 58.7</div> <div style="border: 1px solid black; padding: 2px;">29 Cu copper 63.5</div> <div style="border: 1px solid black; padding: 2px;">30 Zn zinc 65.4</div> <div style="border: 1px solid black; padding: 2px;">31 Ga gallium 69.7</div> <div style="border: 1px solid black; padding: 2px;">32 Ge germanium 72.6</div> <div style="border: 1px solid black; padding: 2px;">33 As arsenic 74.9</div> <div style="border: 1px solid black; padding: 2px;">34 Se selenium 79.0</div> <div style="border: 1px solid black; padding: 2px;">35 Br bromine 79.9</div> <div style="border: 1px solid black; padding: 2px;">36 Kr krypton 83.8</div> </div>																	
		<div style="display: flex; justify-content: space-between;"> <div style="border: 1px solid black; padding: 2px;">37 Rb rubidium 85.5</div> <div style="border: 1px solid black; padding: 2px;">38 Sr strontium 87.6</div> </div>																	
		<div style="display: flex; justify-content: space-between;"> <div style="border: 1px solid black; padding: 2px;">39 Y yttrium 88.9</div> <div style="border: 1px solid black; padding: 2px;">40 Zr zirconium 91.2</div> <div style="border: 1px solid black; padding: 2px;">41 Nb niobium 92.9</div> <div style="border: 1px solid black; padding: 2px;">42 Mo molybdenum 95.9</div> <div style="border: 1px solid black; padding: 2px;">43 Tc technetium —</div> <div style="border: 1px solid black; padding: 2px;">44 Ru ruthenium 101.1</div> <div style="border: 1px solid black; padding: 2px;">45 Rh rhodium 102.9</div> <div style="border: 1px solid black; padding: 2px;">46 Pd palladium 106.4</div> <div style="border: 1px solid black; padding: 2px;">47 Ag silver 107.9</div> <div style="border: 1px solid black; padding: 2px;">48 Cd cadmium 112.4</div> <div style="border: 1px solid black; padding: 2px;">49 In indium 114.8</div> <div style="border: 1px solid black; padding: 2px;">50 Sn tin 118.7</div> <div style="border: 1px solid black; padding: 2px;">51 Sb antimony 121.8</div> <div style="border: 1px solid black; padding: 2px;">52 Te tellurium 127.6</div> <div style="border: 1px solid black; padding: 2px;">53 I iodine 126.9</div> <div style="border: 1px solid black; padding: 2px;">54 Xe xenon 131.3</div> </div>																	
		<div style="display: flex; justify-content: space-between;"> <div style="border: 1px solid black; padding: 2px;">55 Cs caesium 132.9</div> <div style="border: 1px solid black; padding: 2px;">56 Ba barium 137.3</div> </div>																	
		<div style="display: flex; justify-content: space-between;"> <div style="border: 1px solid black; padding: 2px;">57–71 lanthanoids</div> <div style="border: 1px solid black; padding: 2px;">72 Hf hafnium 178.5</div> <div style="border: 1px solid black; padding: 2px;">73 Ta tantalum 180.9</div> <div style="border: 1px solid black; padding: 2px;">74 W tungsten 183.8</div> <div style="border: 1px solid black; padding: 2px;">75 Re rhenium 186.2</div> <div style="border: 1px solid black; padding: 2px;">76 Os osmium 190.2</div> <div style="border: 1px solid black; padding: 2px;">77 Ir iridium 192.2</div> <div style="border: 1px solid black; padding: 2px;">78 Pt platinum 195.1</div> <div style="border: 1px solid black; padding: 2px;">79 Au gold 197.0</div> <div style="border: 1px solid black; padding: 2px;">80 Hg mercury 200.6</div> <div style="border: 1px solid black; padding: 2px;">81 Tl thallium 204.4</div> <div style="border: 1px solid black; padding: 2px;">82 Pb lead 207.2</div> <div style="border: 1px solid black; padding: 2px;">83 Bi bismuth 209.0</div> <div style="border: 1px solid black; padding: 2px;">84 Po polonium —</div> <div style="border: 1px solid black; padding: 2px;">85 At astatine —</div> <div style="border: 1px solid black; padding: 2px;">86 Rn radon —</div> </div>																	
		<div style="display: flex; justify-content: space-between;"> <div style="border: 1px solid black; padding: 2px;">87 Fr francium —</div> <div style="border: 1px solid black; padding: 2px;">88 Ra radium —</div> </div>																	
		<div style="display: flex; justify-content: space-between;"> <div style="border: 1px solid black; padding: 2px;">89–103 actinoids</div> <div style="border: 1px solid black; padding: 2px;">104 Rf rutherfordium —</div> <div style="border: 1px solid black; padding: 2px;">105 Db dubnium —</div> <div style="border: 1px solid black; padding: 2px;">106 Sg seaborgium —</div> <div style="border: 1px solid black; padding: 2px;">107 Bh bohrium —</div> <div style="border: 1px solid black; padding: 2px;">108 Hs hassium —</div> <div style="border: 1px solid black; padding: 2px;">109 Mt meitnerium —</div> <div style="border: 1px solid black; padding: 2px;">110 Ds darmstadtium —</div> <div style="border: 1px solid black; padding: 2px;">111 Rg roentgenium —</div> <div style="border: 1px solid black; padding: 2px;">112 Cn copernicium —</div> <div style="border: 1px solid black; padding: 2px;">113 Nh nihonium —</div> <div style="border: 1px solid black; padding: 2px;">114 Fl flerovium —</div> <div style="border: 1px solid black; padding: 2px;">115 Mc moscovium —</div> <div style="border: 1px solid black; padding: 2px;">116 Lv livermorium —</div> <div style="border: 1px solid black; padding: 2px;">117 Ts tennessine —</div> <div style="border: 1px solid black; padding: 2px;">118 Og oganeson —</div> </div>																	

lanthanoids

actinoids

57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.4	61 Pm promethium —	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1	71 Lu lutetium 175.0
89 Ac actinium	90 Th thorium 232.0	91 Pa protactinium 231.0	92 U uranium 238.0	93 Np neptunium —	94 Pu plutonium —	95 Am americium —	96 Cm curium —	97 Bk berkelium —	98 Cf californium —	99 Es einsteinium —	100 Fm fermium —	101 Md mendelevium —	102 No nobelium —	103 Lr lawrencium —