Cambridge IGCSE[™] Additional Mathematics (0606) and Cambridge O Level Additional Mathematics (4037)

List of formulas

Equation of a circle with centre (a, b) and radius r.

$$(x-a)^2 + (y-b)^2 = r^2$$

Curved surface area, A, of cone of radius r, sloping edge l.

 $A = \pi r l$

Surface area, A, of sphere of radius r.

$$A = 4\pi r^2$$

Volume, *V*, of pyramid or cone, base area *A*, height *h*.

$$V = \frac{1}{3}Ah$$

Volume, V, of sphere of radius r.

$$V = \frac{4}{3}\pi r^3$$

Quadratic equation

For the equation
$$ax^2 + bx + c = 0$$
,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial theorem

$$(a+b)^{n} = a^{n} + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^{2} + \dots + \binom{n}{r}a^{n-r}b^{r} + \dots + b^{n},$$

where *n* is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

Arithmetic series

$$u_n = a + (n-1)d$$

 $S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a + (n-1)d\}$

Geometric series

$$u_n = ar^{n-1}$$

$$S_n = \frac{a(1-r^n)}{1-r} \quad (r \neq 1)$$

$$S_{\infty} = \frac{a}{1-r} \quad (|r| < 1)$$

Identities

$$\sin^2 A + \cos^2 A = 1$$
$$\sec^2 A = 1 + \tan^2 A$$
$$\csc^2 A = 1 + \cot^2 A$$

Formulas for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$
$$a^2 = b^2 + c^2 - 2bc \cos A$$
$$\Delta = \frac{1}{2} ab \sin C$$

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (Cambridge University Press & Assessment) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge Assessment International Education is part of Cambridge University Press & Assessment. Cambridge University Press & Assessment is a department of the University of Cambridge.