



Cambridge O Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



ADDITIONAL MATHEMATICS

4037/01

Paper 1 Non-calculator

For examination from 2025

PRACTICE PAPER

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- Calculators must **not** be used in this paper.
- You must show all necessary working clearly.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages.

List of formulas

Equation of a circle with centre (a, b) and radius r . $(x - a)^2 + (y - b)^2 = r^2$

Curved surface area, A , of cone of radius r , sloping edge l . $A = \pi rl$

Surface area, A , of sphere of radius r . $A = 4\pi r^2$

Volume, V , of pyramid or cone, base area A , height h . $V = \frac{1}{3}Ah$

Volume, V , of sphere of radius r . $V = \frac{4}{3}\pi r^3$

Quadratic equation For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial theorem $(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n$,

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

Arithmetic series $u_n = a + (n - 1)d$
 $S_n = \frac{1}{2}n(a + l) = \frac{1}{2}n\{2a + (n - 1)d\}$

Geometric series $u_n = ar^{n-1}$
 $S_n = \frac{a(1 - r^n)}{1 - r} \quad (r \neq 1)$
 $S_\infty = \frac{a}{1 - r} \quad (|r| < 1)$


Identities $\sin^2 A + \cos^2 A = 1$
 $\sec^2 A = 1 + \tan^2 A$
 $\operatorname{cosec}^2 A = 1 + \cot^2 A$

Formulas for $\triangle ABC$ $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$
 $a^2 = b^2 + c^2 - 2bc \cos A$
 $\Delta = \frac{1}{2} ab \sin C$

- 1 The curve $y = 2x^2 + k + 4$ intersects the straight line $y = (k + 4)x$ at two distinct points. Find the possible values of k .

[4]

2



The diagram shows the graph of $y = f(x)$, where $f(x)$ is a cubic polynomial.

(a) Find $f(x)$ giving your answer as a product of linear factors. [3]

(b) Write down the values of x such that $f(x) < 0$. [2]


3 Solve the inequality $|5x + 4| \leq |2x - 3|$. [4]

4 The function f is defined as $f(x) = x^2 + 2x - 3$ for $x \geq -1$.

(a) Given that the minimum value of $x^2 + 2x - 3$ occurs when $x = -1$ explain why $f(x)$ has an inverse. [1]

(b) On the axes below, sketch the graph of $y = f(x)$ and the graph of $y = f^{-1}(x)$.

Label each graph and state the intercepts on the coordinate axes. [4]



5 (a) Show that $\frac{\sin x \tan x}{1 - \cos x}$ can be written as $1 + \sec x$. [4]

(b) Hence show that $\frac{d}{dx} \left(\frac{\sin x \tan x}{1 - \cos x} \right)$ can be written as $\tan x \sec x$. [2]

6 Solve the following simultaneous equations.

$$\log_3(x + y) = 2$$

$$2\log_3(x + 1) = \log_3(y + 2)$$

[6]

- 7 When y^3 is plotted against $\ln x$, a straight-line graph is obtained. The straight line passes through the points (1, 5) and (6, 15).

(a) Find y in terms of x .

[4]

(b) Given that $y > -1$, find the corresponding values of x .

[2]

8 The polynomial $p(x) = 6x^3 + ax^2 + bx + 2$ where a and b are integers, has a factor of $x - 2$.

(a) Given that $p(1) = -2p(0)$, find the value of a and of b . [4]

(b) Using your values of a and b ,

(i) find the remainder when $p(x)$ is divided by $2x - 1$, [2]

(ii) factorise $p(x)$. [2]


- 9 (a) Find the equation of the tangent to the curve $y = x^3 - 6x^2 + 3x + 10$ at the point where $x = 1$.
Give your answer in the form $y = mx + c$. [4]

- (b) Find the coordinates of the point where this tangent meets the curve again. [5]

10 Find the exact value of $\int_2^4 \frac{(x+1)^2}{x^2} dx$.

[6]

11



In the diagram $\vec{OP} = 2\mathbf{b}$, $\vec{OS} = 3\mathbf{a}$, $\vec{SR} = \mathbf{b}$ and $\vec{PQ} = \mathbf{a}$.
The lines OR and QS intersect at the point X .

(a) Find \vec{OQ} in terms of \mathbf{a} and \mathbf{b} . [1]

(b) Find \vec{QS} in terms of \mathbf{a} and \mathbf{b} . [1]

(c) Given that $\vec{OX} = \mu\vec{QS}$, find \vec{OX} in terms of \mathbf{a} , \mathbf{b} and μ . [1]

(d) Given that $\vec{OX} = \lambda\vec{OR}$, find \vec{OX} in terms of \mathbf{a} , \mathbf{b} and λ . [1]

(e) Find the values of λ and μ .

[3]

12 Circle C_1 has equation $x^2 + y^2 - 2x + 6y = 0$.

The point A lies on the circumference of the circle and has coordinates $(2, 0)$.

(a) Find the equation of the tangent to the circle at A .

[5]

The circle C_1 is reflected in the tangent at A to form the circle C_2 .

(b) Find the equation of C_2 .

[5]

Question 13 is printed on the next page.

- 13 It is given that $2 + \cos \theta = x$ for $1 < x < 3$ and $2 \operatorname{cosec} \theta = y$ for $y > 2$.
Find y in terms of x .

[4]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (Cambridge University Press & Assessment) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge Assessment International Education is part of Cambridge University Press & Assessment. Cambridge University Press & Assessment is a department of the University of Cambridge.