

Cambridge International AS & A Level

BIOLOGY			9700/23
Paper 2 AS Level St	uctured Questions	Octo	ber/November 2024
MARK SCHEME			
Maximum Mark: 60			
	Publi	shed	

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2024 series for most Cambridge IGCSE, Cambridge International A and AS Level components, and some Cambridge O Level components.

PUBLISHED

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptions for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond
 the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Science-Specific Marking Principles

- 1 Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.
- 2 The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.
- Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).
- The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.

5 <u>'List rule' guidance</u>

For questions that require *n* responses (e.g. State **two** reasons ...):

- The response should be read as continuous prose, even when numbered answer spaces are provided.
- Any response marked *ignore* in the mark scheme should not count towards *n*.
- Incorrect responses should not be awarded credit but will still count towards *n*.
- Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should not be
 awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this
 should be treated as a single incorrect response.
- Non-contradictory responses after the first *n* responses may be ignored even if they include incorrect science.

6 Calculation specific guidance

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form (e.g. $a \times 10^n$) in which the convention of restricting the value of the coefficient (a) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

7 Guidance for chemical equations

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.

PUBLISHED

Mark scheme abbreviations

; separates marking points

I alternative answers for the same point

A accept (for answers correctly cued by the question, or by extra guidance)

R reject ignore

the word / phrase in brackets is not required, but sets the contextalternative wording (where responses vary more than usual)

underline actual word given must be used by candidate (grammatical variants accepted)

max indicates the maximum number of marks that can be given

ora or reverse argument

mp marking point (with relevant number)

ecf error carried forward AVP alternative valid point

Question	Answer	Marks
1(a)(i)	B and C; Accept A, B and C if any other letters do not award the mark, ignore names	1
1(a)(ii)	any two from:	2
	keeps the (named) airway(s) open / prevents collapse of airways; AW R bronchiole / alveoli e.g. to allow free flow of air into and out of, airways / lungs e.g. provides support (for the named airway(s)) I 'provides strength'	
	allows flexibility; A described e.g. bending neck, swallowing food	
	AVP; e.g. (C-shaped incomplete) rings (in, B / trachea) allow, lengthening / widening, during breathing in / inspiration / inhalation	

Question		Answer					
1(a)(iii)	one ma if find a feature	region one mark for each letter – for 1.3 allow two or three letters if all are correct – ignore names one mark for feature matched with function – feature and relationship to function must match if find a correct feature and its function in the last column – accept feature must be visible in Fig. 1.2 and Fig. 1.3 if gas (as in gas exchange) is not used, then answer must refer to oxygen and carbon dioxide					
	Fig.	region of the gas exchange system (A, B, C, D, E, F, G)	one visible feature	one way in which the feature relates to its function			
	1.2	G;	air spaces	large surface area for, diffusion / gas exchange;			
			walls are, thin / one cell thick A squamous epithelium R cell wall	short distance for, diffusion / gas exchange;			
			good blood supply / capillaries A capillary / red blood cells	for, diffusion / gas exchange or maintains steep concentration gradients or absorption / transport, of oxygen;			
			arteriole / venule R artery / vein	supply blood to capillaries / carry blood away from capillaries ;			
	1.3	B; Accept C/A	goblet cells	secrete / produce / make, mucus ;			
			(many) cilia / ciliated epithelial cells / ciliated epithelium / ciliated cells	(cilia) move / AW, mucus (towards back of throat / AW);			
			ciliated epithelium	provides barrier to pathogens;			

Question	Answer	Marks
1(b)	any five from:	5
	sinoatrial node, releases / generates / produces, impulses / waves of excitation / waves of depolarisation / AW; I 'signals' A SAN / SA node / pacemaker R nervous impulses	
	2 impulses / AW, spreads to / across, walls / muscle of, (right) atrium; A atria	
	3 <u>right atrium</u> contracts ; A atrial systole if right atrium is given in answer, A pumps for contracts	
	4 tricuspid / (right) atrioventricular, valve <u>opens</u> to allow blood to flow (from right atrium) into <u>right ventricle</u> ; R bicuspid valve / left atrioventricular valve	
	5 atrioventricular node sends impulses down, septum / Purkyne tissue; A to apex of heart / down Bundle of His A AVN / AV node	
	6 atrioventricular node is responsible for / AW, a time delay; A AVN / AV node	
	7 <u>right ventricle</u> contracts ; A ventricular systole if right ventricle is given in answer, A pumps for contracts	
	8 semi-lunar valve opens and blood flows into pulmonary artery / arteries; A pulmonary valve R aorta	
	9 AVP ; e.g. tricuspid / (right) atrioventricular, valve <u>closes</u> to stop, backflow / blood entering right atrium (only in correct context)	

om: I repair cells place (old / worn-out / dead / short-lived) cells / repair (damaged) tissue; cells divide and, differentiate / become specialised; A form different types of cells of self-renewal e.g. stem cells divide to maintain the, pool / number, of stem cells; increase in number of stem cells ref. to movement of food through intestine / presence of pathogens, increases chance of damage cells (produced by stem cells) are genetically identical so the function of, the intestine / organs / tissues, can continue / AW lea that allows growth (of small intestine) during development / AW R growth of cells	2
cells divide and, differentiate / become specialised; A form different types of cells of self-renewal e.g. stem cells divide to maintain the, pool / number, of stem cells; increase in number of stem cells ref. to movement of food through intestine / presence of pathogens, increases chance of damage cells (produced by stem cells) are genetically identical so the function of, the intestine / organs / tissues, can continue / AW	
of self-renewal e.g. stem cells divide to maintain the, pool / number, of stem cells; increase in number of stem cells ref. to movement of food through intestine / presence of pathogens, increases chance of damage cells (produced by stem cells) are genetically identical so the function of, the intestine / organs / tissues, can continue / AW	
ref. to movement of food through intestine / presence of pathogens, increases chance of damage ells (produced by stem cells) are genetically identical so the function of, the intestine / organs / tissues, can portione / AW	
ells (produced by stem cells) are genetically identical so the function of, the intestine / organs / tissues, can ontinue / AW	
	4
;	1
s;	1
от	2
sister / identical, <u>chromatids</u> attached together ; R 'sister chromosomes' holds the chromatids together after, replication / S phase	
romeres) divide / separate, (at the start) in anaphase ; A end of metaphase	
ughter chromosomes can be moved to expecite poles. A cister/identical chromatide	
n tr	(chromatids / chromosomes) to spindle (fibres); in context of mitosis R if stated in interphase A kinetochore or spindle microtubules for spindle (fibres) tromeres) divide / separate, (at the start) in anaphase; A end of metaphase sughter chromosomes can be moved to opposite poles; A sister / identical, chromatids ward MP4 unless MP3 already given

Question		Answer				
2(c)	for ea	or function in explanation if not on first line ch cell – one mark for the function and or each cell straight through to max 2		6		
	cell	function	explanation			
	Р	absorption / uptake / movement in, of, (named) substances / nutrients	microvilli, to give a large surface area / increase surface area / increase number of transport proteins (for uptake of digested food)			
		diffusion of, (named) substances / nutrients	I cilia have large surface area A large surface area:volume ratio for microvilli			
		active uptake / active transport, of (named) substances / of nutrients	mitochondria to provide, ATP / energy, for, active transport / active uptake R produce / make, energy R energy for cilia			
		breakdown of food molecules	large surface of membrane for enzymes for digestion			
		production / secretion, of enzymes	microvilli for membrane enzymes / RER for making enzymes / Golgi for packaging enzymes <i>or</i> modification of enzymes			
	Q	secretion / release / exocytosis / AW,	rough endoplasmic reticulum for making protein (part of mucin / mucus)			
		of mucin / mucus I produce and provide	Golgi body for, packaging / glycosylating / modification / AW, mucin A mucus AW = adding carbohydrate / sugars			
		if Q is secreting protein / unnamed substance allow ECF for explanation	(large) vesicles / vacuoles, for storage / for containing / for transport, of mucin A mucus			
	R	secretion / release / exocytosis / move ment out / AW, of (named) protein(s)	(large quantity of) RER for, making proteins / making enzymes / translation / AW			
		I produce and provide A hormones / peptides / enzymes	Golgi body for packaging / modification / AW, proteins			
		,	vacuoles / vesicles, for transport to cell surface membrane / release by exocytosis / fusion with cell surface membrane			

Question	Answer	Marks
3(a)(i)	pyrimidine; accept phonetic or close spellings, e.g. pyramidine / pyrimadine / pirimadine / primidine	1
3(a)(ii)	any one from:	1
	each nucleic acid has 4 (types of) base; A has 4 bases DNA has A, T, C and G and RNA has A, U, C and G; A full names even if misspelt unless another molecule (e.g. thiamine) idea that in RNA U replaces T / DNA has T but RNA has U;	
3(b)	 any four from: max 2 if answer is about transcription and mRNA penalise polypeptide once 1 (triphosphate) activated / free, nucleotide (with thymine); 2 thymine / T, pairs with, adenine / A; I complementary base pairing unqualified 3 adenine / A, on template strand; 4 formation of two hydrogen bonds between T and A; I H bonds between C and G 5 ref to DNA polymerase in correct context (e.g. correct base pairing / bond formation / proof reading); 6 forms phosphodiester bond (between -OH of deoxyribose and phosphate of nucleotide); R if DNA ligase 7 release of, pyrophosphate / P-P / two phosphates; 8 nucleotide added to the 3' end of, newly-synthesised strand / leading strand / daughter strand (of DNA); A extension is in the 5' to 3' direction 	4
3(c)	 strands (are parallel and) run in opposite directions / one strand in 5′ to 3′ direction and other strand in 3′ to 5′ direction; any two from: elongation / synthesis / extension / AW, of strand is, in 5′ to 3′ direction / not in 3′ to 5′ direction; DNA polymerase, moves in the 5′ to 3′ direction / can only add nucleotides to the 3′ end; idea that phosphate of each nucleotide is added to C3 of last nucleotide of growing strand; leading strand is made continuously / lagging strand is made up of Okazaki fragments; Okazaki fragments are attached together by DNA ligase; 	3

Question	Answer	Marks
4(a)	any four from: I 'find optimum conditions for the enzyme mixture' A 'enzymes' for enzyme mixture	4
	 1 (find) suitable, raw material / crop wastes / substrate; using Table 4.1 2 (find) crop waste that has high concentration of substrate for enzyme M; ORA 	
	 assume answer is about solution or mixture unless told about individual enzymes (M, N and O) 3 (find), optimum pH / optimum temperature; A reference to, pH / temperature, for maximum activity I find optimum pH / temperature of each enzyme 4 (find) appropriate concentrations of the enzymes in the mixture; I 'proportions' 5 idea of ratio of enzyme mixture to, raw material / substrate / crop waste; 6 (find) the best pre-treatment of the, raw material / substrate / crop waste; 7 (find) if any, inhibitors / cofactors, are present in the solution; 8 (find) if any products are toxic; 9 identify the sugars produced; 10 (find) how long it takes to produce sugars / (find) rate of sugar production; 11 ref to enzyme immobilisation; 	
4(b)	any four from:	4
	 cellulose and, (named) protein / polypeptides, were present (in cell wall); e.g. expansins R if collagen cellulose is broken down (fully) to β-glucose / β-glucose is the monomer of cellulose; A 'made of' protein / polypeptide / peptide, is broken down (fully) to amino acids; A 'made of' idea that partial hydrolysis has occurred to form, peptides / short chains of β-glucose; enzymes include cellulase and protease; breakage of glycosidic and peptide bonds; A glucosidic AVP; e.g. beta 1-4 glycosidic linkage / exopeptidases producing amino acids / endopeptidases producing peptides 	

Question	Answer	Marks
5(a)(i)	U4 – T-helper (lymphocyte / cell) AND V4 – T-killer / T-cytotoxic (lymphocyte / cell); A 'helper and killer' I helping and killing	1
5(a)(ii)	 any four from: U4/helper secretes / produces / releases, cytokines / interleukins; I cell signalling molecules / cytokinins to stimulate, B-lymphocytes / B cells, to, divide / develop into plasma cells; A stimulates humoral response / clonal expansion / development of memory cells stimulate macrophages to, be active / carry out phagocytosis / respond (to presence of pathogens) / be 'angry'; I endocytosis 	4
	 V4/killer produces / secretes / releases, chemicals, to kill / break down, infected cells; A (auto)lysis / apoptosis e.g. perforin / hydrogen peroxide / granzymes / proteases; A hydrolytic enzymes / toxins detail of action of named chemical (see below); AVP; e.g. correct ref to CD4 receptors (on U4/T-helper) / CD8 receptors on (V4/T-killer) perforin – makes pores in cell surface membrane of infected cell for delivery of granzymes hydrogen peroxide – oxidises cell components / stimulates apoptosis granzymes / proteases – break down proteins (granzymes are serine proteases) 	

Question			Answer			Marks
5(b)	 ref to herd immunity / vaccinate children as ref to provide booster accurate record keep start vaccination imm education about, effe A counter, misinformation make sure there is a provide funding for valuate funding for valuate or monitor sumaintain cold chain (truse a thermostable valuate different types) 	ole, globally / to whole popular described; e.g. vaccinate a early as possible; is / use effective vaccine so ing of those vaccinated / find ontact tracing / described; e ediately cases are found / rects of the disease / benefits ation / antivaxers A important sufficient supply of the vaccinacine, production / distributionnel; ing people with the disease uccess of vaccine is effective accine.	d those who have not been e.g. look out for any cases of ef to ring immunity / benefits of having the vaccine / transince of vaccination sine; ion / delivery; A provide, chamme e) (oral) and Salk (injected); d	; e.g. live / attenuated, virus vaccinated; f the disease of ring immunity;	ne	4
6(a)	feature	plant cell	prokaryotic cell	virus		4
	external structure	cell wall composed of cellulose	cell wall composed of peptidoglycan / murein;	capsid composed of protein / polypeptides; A capsomeres		
	size of ribosomes	80S and 70S	70S;	no ribosomes		
	nucleic acids	DNA and RNA	DNA and RNA	DNA or RNA ; R DNA and RNA		

Question	Answer	Marks
6(b)(i)	any one from:	1
	untreated / raw, sewage; A poor sewage treatment / poor sewer system contaminated drinking water; contaminated food / crops fertilised by raw sewage / seafood contaminated by raw sewage; AVP; e.g. newly arrived people infected with, <i>Vibrio cholerae</i> / cholera pathogen (e.g. UN troops in Haiti in 2010)	
6(b)(ii)	choleragen is composed of more than one polypeptide; A 6 / 7, polypeptides A amino acid chain A composed of, multiple / several, polypeptides	1
6(b)(iii)	glycolipid / glycoprotein;	1
6(b)(iv)	ATP / adenosine triphosphate ; I adenine triphosphate	1
6(b)(v)	chloride ions decrease the <u>water potential</u> of, intestine lumen / outside of cells / AW; A low <u>er</u> water potential A loss of chloride ions makes the, cytoplasm / the cells, have a less negative / higher, water potential (so water, leaves cell / enters lumen) by osmosis / down water potential gradient / from high to low water potential;	2