

# Cambridge International AS & A Level

---

**BIOLOGY**

**9700/43**

Paper 4 A Level Structured Questions

**October/November 2024**

MARK SCHEME

Maximum Mark: 100

---

**Published**

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2024 series for most Cambridge IGCSE, Cambridge International A and AS Level components, and some Cambridge O Level components.

---

This document consists of **15** printed pages.

**PUBLISHED****Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptions for a question. Each question paper and mark scheme will also comply with these marking principles.

**GENERIC MARKING PRINCIPLE 1:**

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

**GENERIC MARKING PRINCIPLE 2:**

Marks awarded are always **whole marks** (not half marks, or other fractions).

**GENERIC MARKING PRINCIPLE 3:**

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

**GENERIC MARKING PRINCIPLE 4:**

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

**GENERIC MARKING PRINCIPLE 5:**

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

**GENERIC MARKING PRINCIPLE 6:**

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

**Science-Specific Marking Principles**

1 Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.

2 The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.

3 Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).

4 The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.

5 'List rule' guidance

For questions that require *n* responses (e.g. State **two** reasons ...):

- The response should be read as continuous prose, even when numbered answer spaces are provided.
- Any response marked *ignore* in the mark scheme should not count towards *n*.
- Incorrect responses should not be awarded credit but will still count towards *n*.
- Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should **not** be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response.
- Non-contradictory responses after the first *n* responses may be ignored even if they include incorrect science.

**6** Calculation specific guidance

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form (e.g.  $a \times 10^n$ ) in which the convention of restricting the value of the coefficient ( $a$ ) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

**7** Guidance for chemical equations

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.

**Mark scheme abbreviations**

<b>;</b>	separates marking points
<b>/</b>	alternative answers for the same point
<b>A</b>	accept (for answers correctly cued by the question, or by extra guidance)
<b>R</b>	reject
<b>I</b>	ignore
<b>( )</b>	the word / phrase in brackets is not required, but sets the context
<b>AW</b>	alternative wording (where responses vary more than usual)
<b>underline</b>	actual word given must be used by candidate (grammatical variants accepted)
<b>max</b>	indicates the maximum number of marks that can be given
<b>ora</b>	or reverse argument
<b>mp</b>	marking point (with relevant number)
<b>ecf</b>	error carried forward
<b>AVP</b>	alternative valid point

**PUBLISHED**

<b>Question</b>	<b>Answer</b>	<b>Marks</b>
1(a)(i)	<b>A</b> = dendrite(s) ; <b>B</b> = nuclei ; <b>C</b> = synaptic knob(s) ;	<b>3</b>
1(a)(ii)	<b>1</b> rat has, myelin sheath / Schwann cell / myelinated axon / myelinated neurone ;  <b>2</b> (rat has) saltatory conduction <b>OR</b> (rat) impulse / action potential, jumps / leaps / AW, from (one) <u>node of Ranvier</u> to, the next / another ;	<b>2</b>
1(b)	<i>any <b>three</b> from:</i>  <b>snail / Fig. 1.3</b> <b>1</b> greater depolarisation <b>OR</b> (membrane potential) gets more positive / has higher increase / has higher peak ;  <b>2</b> action potential, slower / takes longer / takes more time <b>OR</b> longer time to return to resting potential ;  <b>3</b> longer (absolute / relative) <u>refractory period</u> ;  <b>4</b> longer hyperpolarisation ;  <b>5</b> AVP ;	<b>3</b>

**PUBLISHED**

Question	Answer	Marks
2(a)	<p>any <b>three</b> from:</p> <ol style="list-style-type: none"> <li>1 open in, day / light ;</li> <li>2 to take in carbon dioxide for, photosynthesis / light-independent stage / Calvin cycle ;</li> <li>3 close in, night / dark ;</li> <li>4 to decrease water loss by transpiration ;</li> </ol>	<b>3</b>
2(b)	<p>any <b>six</b> from:</p> <ol style="list-style-type: none"> <li>1 hydrogen ions / protons / H ions / H<sup>+</sup>, leave (cell) using, energy / ATP ;</li> <li>2 low H<sup>+</sup> (in cell) / <u>more</u> negative charge ;</li> <li>3 K<sup>+</sup> / potassium ion, channel (proteins) open <b>OR</b> K<sup>+</sup>, move into / enter, cell (by facilitated diffusion) ;</li> <li>4 Cl<sup>-</sup> ions, move into / enter (cell) ;</li> <li>5 water potential of cell decreases ;</li> <li>6 water moves in (to cell) by <u>osmosis</u> ;</li> <li>7 (cells) become turgid / swell / increase in volume / expand ;</li> <li>8 (guard cell) inner wall thicker / outer wall thinner, so cells, bend / curve ;</li> </ol>	<b>6</b>
2(c)	<ol style="list-style-type: none"> <li>1 (environment) light needed, to keep cycle normal / to achieve same peak % of stomata opening ;</li> <li>2 genes have a role as (regular), rhythm / pattern / cycle, continues, in darkness / without light ;</li> </ol>	<b>2</b>

Question	Answer						Marks																																			
3(a)	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th data-bbox="338 217 943 284" rowspan="2">event</th> <th colspan="4" data-bbox="943 217 1234 284">stage</th> <th data-bbox="1234 217 1279 284"></th> </tr> <tr> <th data-bbox="943 284 1016 347">G</th> <th data-bbox="1016 284 1090 347">LR</th> <th data-bbox="1090 284 1164 347">KC</th> <th data-bbox="1164 284 1234 347">OP</th> <th data-bbox="1234 284 1279 347"></th> </tr> </thead> <tbody> <tr> <td data-bbox="338 347 943 411">catabolism</td> <td data-bbox="943 347 1016 411">✓</td> <td data-bbox="1016 347 1090 411">✓</td> <td data-bbox="1090 347 1164 411">✓</td> <td data-bbox="1164 347 1234 411">✗</td> <td data-bbox="1234 347 1279 411">;</td> </tr> <tr> <td data-bbox="338 411 943 475">coenzyme is reduced or oxidised</td> <td data-bbox="943 411 1016 475">✓</td> <td data-bbox="1016 411 1090 475">✓</td> <td data-bbox="1090 411 1164 475">✓</td> <td data-bbox="1164 411 1234 475">✓</td> <td data-bbox="1234 411 1279 475">;</td> </tr> <tr> <td data-bbox="338 475 943 579">a coenzyme forms a covalent bond with a respiratory intermediate</td> <td data-bbox="943 475 1016 579">✗</td> <td data-bbox="1016 475 1090 579">✓</td> <td data-bbox="1090 475 1164 579">✗</td> <td data-bbox="1164 475 1234 579">✗</td> <td data-bbox="1234 475 1279 579">;</td> </tr> <tr> <td data-bbox="338 579 943 643">carbon dioxide is released</td> <td data-bbox="943 579 1016 643">✗</td> <td data-bbox="1016 579 1090 643">✓</td> <td data-bbox="1090 579 1164 643">✓</td> <td data-bbox="1164 579 1234 643">✗</td> <td data-bbox="1234 579 1279 643">;</td> </tr> </tbody> </table>						event	stage					G	LR	KC	OP		catabolism	✓	✓	✓	✗	;	coenzyme is reduced or oxidised	✓	✓	✓	✓	;	a coenzyme forms a covalent bond with a respiratory intermediate	✗	✓	✗	✗	;	carbon dioxide is released	✗	✓	✓	✗	;	<b>4</b>
event	stage																																									
	G	LR	KC	OP																																						
catabolism	✓	✓	✓	✗	;																																					
coenzyme is reduced or oxidised	✓	✓	✓	✓	;																																					
a coenzyme forms a covalent bond with a respiratory intermediate	✗	✓	✗	✗	;																																					
carbon dioxide is released	✗	✓	✓	✗	;																																					
3(b)(i)	<p><i>any <b>four</b> from:</i></p> <ol style="list-style-type: none"> <li><b>1</b> (RQ =) ratio of carbon dioxide (exhaled) to oxygen (inhaled) ;</li> <li><b>2</b> air flow meter measures oxygen inhaled ;</li> <li><b>3</b> carbon dioxide sensor measures concentration of CO<sub>2</sub> exhaled ;</li> <li><b>4</b> carbohydrate RQ = 1 ;</li> <li><b>5</b> lipid RQ = 0.7 ;</li> </ol>						<b>4</b>																																			
3(b)(ii)	<p><i>any <b>three</b> from:</i></p> <ol style="list-style-type: none"> <li><b>1</b> lipid has higher energy (value) <b>OR</b> carbohydrate 15–17 kJ g<sup>-1</sup> and lipid 37–40 kJ g<sup>-1</sup> ;</li> <li><b>2</b> more, hydrogen (atoms) / C-H bonds, in lipid ;</li> <li><b>3</b> more reduced, NAD / FAD ;</li> <li><b>4</b> greater proton gradient / more protons in intermembrane space / more protons pass through ATP synth(et)ase ;</li> </ol>						<b>3</b>																																			



**PUBLISHED**

Question	Answer	Marks
4(a)	<p><i>any <b>three</b> from:</i> <i>in Venus fly traps</i></p> <ol style="list-style-type: none"> <li>1 touch two (sensory hairs) ;</li> <li>2 protons, leave hinge cells / enter hinge cell walls ;</li> <li>3 Ca<sup>2+</sup> ions move into (hinge) cells ;</li> <li>4 water moves into (hinge) cells ;</li> <li>5 (hinge) cells, become turgid / swell / increase in volume / expand ;</li> </ol>	<b>3</b>
4(b)	<p><i>any <b>two</b> from:</i></p> <ol style="list-style-type: none"> <li>1 less / decreased, surface area (of leaf exposed) ;</li> <li>2 less light absorbed by, chloroplasts / thylakoid membranes / grana / pigments / chlorophyll / light harvesting complexes / photosystems ;</li> <li>3 fewer stomata, exposed / available (to air) ;</li> <li>4 less carbon dioxide, enters / absorbed ;</li> </ol>	<b>2</b>
4(c)	<p><i>any <b>six</b> from:</i></p> <p><i>similarities</i></p> <ol style="list-style-type: none"> <li>1 photoactivation of chlorophyll ;</li> <li>2 (energetic) electrons move, along / down, ETC ;</li> <li>3 chemiosmosis ;</li> <li>4 ATP produced ;</li> </ol> <p><i>differences</i></p> <ol style="list-style-type: none"> <li>5 PSI and PSII in non-cyclic <b>and</b> only PSI in cyclic ;</li> <li>6 photolysis / oxygen produced, in non-cyclic only ; ORA</li> <li>7 reduced NADP in non-cyclic only ; ORA</li> <li>8 electrons return to, same photosystem / PSI, in cyclic ;</li> </ol>	<b>6</b>

**PUBLISHED**

Question	Answer	Marks
5(a)	<p>any <b>two</b> from:</p> <ol style="list-style-type: none"> <li>1 one / single, set of chromosomes ;</li> <li>2 have <b>n</b> (number of) chromosomes / half of diploid number ;</li> <li>3 chromosomes not in homologous pairs / only one of each homologous pair ;</li> <li>4 each chromosome is different in, size / shape / genes / loci ;</li> </ol>	<b>2</b>
5(b)	<p>sexual reproduction: <b>B</b> and <b>D</b> ;  mitosis: two from <b>A, B, C, D</b> ;  meiosis: <b>D</b> ;</p>	<b>3</b>
5(c)	<p>any <b>three</b> from:</p> <ol style="list-style-type: none"> <li>1 to maintain, chromosome / diploid, number ;</li> <li>2 from parents to offspring / from generation to generation ;</li> <li>3 to give genetic variation in, gametes / offspring / horses ;</li> <li>4 too many / extra, sets of chromosomes cause (named) problems ;</li> </ol>	<b>3</b>


Question	Answer	Marks
6(a)(i)	<ol style="list-style-type: none"> <li>1 (excess) amino acids, deaminated / have amino group removed ;</li> <li>2 in the liver ;</li> </ol>	<b>2</b>
6(a)(ii)	<p>any <b>three</b> from:</p> <ol style="list-style-type: none"> <li>1 osmoreceptors in hypothalamus (detect change / are receptors) ;</li> <li>2 effector(s): collecting duct / distal convoluted tubule / nephron ;</li> <li>3 target cells: collecting duct (cells) ;</li> </ol>	<b>3</b>

**PUBLISHED**

Question	Answer	Marks
6(b)	<p>any <b>three</b> from:</p> <ol style="list-style-type: none"> <li>1 ultrafiltration / forms glomerular filtrate ;</li> <li>2 high, hydrostatic / blood, <u>pressure</u> in, glomerulus / capillaries ;</li> <li>3 so, water / (named) solutes, move (out of glomerulus / into Bowman's capsule) ;</li> <li>4 through <i>two of</i>: fenestrae / fenestrations basement membrane slit pores / filtration slits / between podocytes ;</li> <li>5 AVP ;</li> </ol>	<b>3</b>
6(c)	<p>any <b>three</b> from:</p> <ol style="list-style-type: none"> <li>1 high(er / est), U:P / ratio → can tolerate water shortage ;</li> <li>2 high(er / est), U:P / ratio → much water reabsorbed / small volume of urine produced ;</li> <li>3 high(er / est), U:P / ratio → can, concentrate urine / conserve water <b>OR</b> high(er / est), U:P / ratio → (well) adapted to dry environment ;</li> <li>4 AVP ;</li> </ol>	<b>3</b>

Question	Answer	Marks
7(a)(i)	<p>any <b>three</b> from:</p> <ol style="list-style-type: none"> <li>1 disruptive / diversifying (selection) ;</li> <li>2 extremes / carnivores and detritus feeders, survive / are selected for <b>and</b> intermediates, die / are selected against ;</li> <li>3 high / intense, competition ;</li> <li>4 intermediates, outcompeted for / don't get enough, detritus / algae, and, (fairy) shrimp / small animals ;</li> <li>5 lack of / limited, food is selection pressure ;</li> <li>6 AVP ;</li> </ol>	<b>3</b>

**PUBLISHED**

Question	Answer	Marks
7(a)(ii)	<p>any <b>three</b> from:</p> <ol style="list-style-type: none"> <li>1 <u>stabilising</u> (selection) ;</li> <li>2 extremes / carnivores and detritus feeders, die / are selected against / are not selected for, <b>and</b> intermediates, survive / are selected for ;</li> <li>3 low / less / prevents, competition ;</li> <li>4 intermediates can eat, both types / all / a wider range, of food ;</li> <li>5 greater variety in the diet improves the growth and development of (intermediate) tadpoles ;</li> </ol>	<b>3</b>
7(a)(iii)	<div style="text-align: center;">  </div> <p>normal distribution curve ;</p>	<b>1</b>
7(a)(iv)	<p>any <b>two</b> from:</p> <ol style="list-style-type: none"> <li>1 new species form due to reproductive isolation ;</li> <li>2 caused by, ecological / behavioural, separation / isolation / differences ;</li> <li>3 in same <u>geographical</u> region ;</li> <li>4 AVP ;</li> </ol>	<b>2</b>
7(b)	<p>any <b>three</b> from:</p> <ol style="list-style-type: none"> <li>1 <i>ref. to</i> DNA sequence (for all / three species) ;</li> <li>2 find / count, nucleotide / base, differences / similarities ;</li> <li>3 fewer differences means, more closely related / less time since divergence / less time to common ancestor ;</li> <li>4 bioinformatics / database / software / BLAST ;</li> <li>5 <i>S. hammondii</i> and <i>S. bombifrons</i>, have the fewest genetic differences / are the most genetically similar ;</li> </ol>	<b>3</b>

**PUBLISHED**

Question	Answer	Marks																
8(a)	<p>any <b>three</b> from:</p> <ol style="list-style-type: none"> <li>1 make cDNA from, messenger RNA / mRNA ;</li> <li>2 using reverse transcriptase ;</li> <li>3 synthesise (gene chemically) from <u>nucleotides</u> / join <u>nucleotides</u> ;</li> <li>4 (using known) amino acid / nucleotide, sequence from database ;</li> </ol>	<b>3</b>																
8(b)	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 25%;">role in genetic engineering</th> <th style="width: 25%;">DNA ligase</th> <th style="width: 25%;">DNA polymerase</th> <th style="width: 25%;"></th> </tr> </thead> <tbody> <tr> <td>joins two sections of sugar phosphate backbone in DNA</td> <td style="text-align: center;">✓</td> <td style="text-align: center;">x</td> <td style="text-align: center;">;</td> </tr> <tr> <td>adds a gene to a plasmid</td> <td style="text-align: center;">✓</td> <td style="text-align: center;">x</td> <td style="text-align: center;">;</td> </tr> <tr> <td>adds free activated DNA nucleotides to a polynucleotide</td> <td style="text-align: center;">x</td> <td style="text-align: center;">✓</td> <td style="text-align: center;">;</td> </tr> </tbody> </table>	role in genetic engineering	DNA ligase	DNA polymerase		joins two sections of sugar phosphate backbone in DNA	✓	x	;	adds a gene to a plasmid	✓	x	;	adds free activated DNA nucleotides to a polynucleotide	x	✓	;	<b>3</b>
role in genetic engineering	DNA ligase	DNA polymerase																
joins two sections of sugar phosphate backbone in DNA	✓	x	;															
adds a gene to a plasmid	✓	x	;															
adds free activated DNA nucleotides to a polynucleotide	x	✓	;															
8(c)	<ol style="list-style-type: none"> <li>1 90–98°C – DNA, denatures / strands separate ;</li> <li>2 50–65°C – primers, bind / base pair / anneal (to DNA) ;</li> <li>3 68–75°C – Taq / DNA, polymerase, makes, DNA / (new) strand ;</li> </ol>	<b>3</b>																

**PUBLISHED**

Question	Answer	Marks
9(a)	<p>any <b>four</b> from:</p> <p>1 both, <u>eukaryotic</u> / <u>eukaryotes</u> / in domain <u>Eukarya</u> ;</p> <p>2 example of eukaryotic feature ;</p> <p><i>Protoctista</i></p> <p>3 example of way in which they vary (<i>must give both</i>) ;</p> <p><i>Fungi (max 3)</i></p> <p>4 cell wall of, chitin / mannan / glucan ;</p> <p>5 hyphae / mycelium ;</p> <p>6 multinucleate / syncytium / many nuclei per cell ;</p> <p>7 heterotrophic / saprotrophic / parasitic ;</p> <p>8 store glycogen ;</p> <p>9 reproduce by spores ;</p>	<b>4</b>
9(b)(i)	<p>any <b>three</b> from:</p> <p>1 measures / gives information about, air, quality / pollution ;</p> <p>2 other species may be harmed by, sulfur dioxide / acid rain / decrease in lichen (abundance / diversity) ;</p> <p>3 <i>ref. to</i> food web / species that feed on lichens ;</p> <p>4 AVP ;</p>	<b>3</b>
9(b)(ii)	<p>any <b>two</b> from:</p> <p>1 role in food web / eaten by other (named) species ;</p> <p>2 provide, shelter / camouflage (for insects / invertebrates) ;</p> <p>3 clean the air / remove (named) pollutants / absorb toxins ;</p> <p>4 may have medical use ;</p> <p>5 <i>ref. to</i> other practical use ;</p> <p>6 primary colonisers / pioneer species / soil formation ;</p> <p>7 ethical / moral / aesthetic, reason ;</p> <p>8 to conserve genetic diversity / future use for genes ;</p>	<b>2</b>

Question	Answer	Marks									
10(a)	(melanic x melanic) $Aa \times Aa$ (Gametes) $A \ a$ and $A \ a$ ;  (Offspring) $AA \ Aa \ Aa \ aa$ ; (melanic, melanic, melanic, typical)  (Ratio) 3 melanic : 1 typical ;	3									
10(b)	it / (melanic) allele, is dominant ;	1									
10(c)(i)	melanic / dominant phenotype, is crossed with homozygous recessive ;	1									
10(c)(ii)	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 30%;"></th> <th style="width: 35%;">same locus (A / a)</th> <th style="width: 35%;">different loci (A / a and B / b)</th> </tr> </thead> <tbody> <tr> <td>genotypes of <b>melanic</b> moths from cross 1</td> <td>AA Aa (aA) ;</td> <td>AaBb Aabb aaBb ;</td> </tr> <tr> <td>proportion of test crosses (cross 2) giving 100% melanic offspring</td> <td>1 in 3</td> <td>none / 0 ;</td> </tr> </tbody> </table>		same locus (A / a)	different loci (A / a and B / b)	genotypes of <b>melanic</b> moths from cross 1	AA Aa (aA) ;	AaBb Aabb aaBb ;	proportion of test crosses (cross 2) giving 100% melanic offspring	1 in 3	none / 0 ;	3
	same locus (A / a)	different loci (A / a and B / b)									
genotypes of <b>melanic</b> moths from cross 1	AA Aa (aA) ;	AaBb Aabb aaBb ;									
proportion of test crosses (cross 2) giving 100% melanic offspring	1 in 3	none / 0 ;									
10(d)	$(24 \times 29) \div 8$ ; $= 87$ ;	2									